

Geologica Macedonica

Journal of the Geological Institute at the Faculty of Natural and
Technical Sciences, University "Goce Delčev" - Štip, R. Macedonia

Kutnahorite - Sasa



<i>Geologica Macedonica</i>	Vol.	No	pp.	Štip
<i>Geologica Macedonica</i>	Год. 34	Број 1	1-84	ШТИП 2020
			стр.	

<i>Geologica Macedonica</i>	Vol.	34	No	1	pp.	1–84	Štip	2020
<i>Geologica Macedonica</i>	Год.		Број		стр.		Штип	

GEOLOGICA MACEDONICA

Published by: – Издава:

"Goce Delčev" University in Štip, Faculty of Natural and Technical Sciences, Štip, North Macedonia
Универзитет „Гоце Делчев“ во Штип, Факултет за природни и технички науки, Штип, Северна Македонија

EDITORIAL BOARD

Todor Serafimovski (N. Macedonia, *Editor in Chief*), **Blažo Boev** (N. Macedonia, *Editor*),
David Alderton (UK), Tadej Dolenc (Slovenia), Ivan Zagorchev (Bulgaria), Wolfgang Todt (Germany),
Nikolay S. Bortnikov (Russia), Clark Burchfiel (USA), Thierry Augé (France), Todor Delipetrov (N. Macedonia),
Vlado Bermanec (Croatia), Milorad Jovanovski (N. Macedonia), Spomenko Mihajlović (Serbia), Dragan Milovanović
(Serbia), Dejan Prelević (Germany), Albrecht von Quadt (Switzerland), Sabina Strmić-Palinkaš (Norway).

УРЕДУВАЧКИ ОДБОР

Тодор Серафимовски (С. Македонија, *главен уредник*), **Блажо Боев** (С. Македонија, *уредник*),
Дејвид Олдертон (В. Британија), Тадеј Доленец (Словенија), Иван Загорчев (Бугарија), Волфганг Тод
(Германија), Николај С. Бортников (Русија), Кларк Барчфил (САД), Тиери Оже (Франција),
Тодор Делипетров (С. Македонија), Владо Берманец (Хрватска), Милорад Јовановски (С. Македонија),
Споменко Михајловиќ (Србија), Драган Миловановиќ (Србија), Дејан Прелевиќ (Германија),
Албрехт фон Квад (Швајцарија), Сабина Стрмиќ-Палинкаш (Норвешка)

Language editor Лектура

Proof-Reading-Service.com, Ltd, Devonshire
Business Centre, Works Road, Letchworth Garden City,
Hertfordshire, SG6 1GJ, United Kingdom
(English) (англиски)

Georgi Georgievski **Георги Георгиевски**
(Macedonian) (македонски)

Technical editor Технички уредник
Blagoja Bogatinoski **Благоја Богатиноски**

Proof-reader Коректор
Alena Georgievska **Алена Георгиевска**

Address	Адреса
GEOLOGICA MACEDONICA	GEOLOGICA MACEDONICA
EDITORIAL BOARD	РЕДАКЦИЈА
Faculty of Natural and Technical Sciences	Факултет за природни и технички науки
P. O. Box 96	пошт. факс 96
MK-2000 Štip, North Macedonia	MK-2000 Штип, Северна Македонија
Tel. ++ 389 032 550 575	Тел. 032 550 575
E-mail: todor.serafimovski@ugd.edu.mk	

400 copies Тираж: 400
Published twice yearly Излегува два пати годишно

Printed by: Печати:
2nd August – Štip 2nd Август – Штип

Price: 10 € Цена: 500 ден.
The edition is published in June 2020 Бројот е отпечатен во јуни 2020,

<i>Geologica Macedonica</i>	Vol.	No	pp.	Štip
<i>Geologica Macedonica</i>	Год.	Број	стр.	Штип
	34	1	1–84	2020

TABLE OF CONTENTS

356. Miha Bukleski

Infra red specular reflectance spectra of pressed polycrystalline samples of alums –
comparison with single crystal spectra 5–14

357. Tena Šijakova-Ivanova, Blažo Boev, Lidija Robeva Čukovska

Preliminary investigations of goethite and monazite from Sivec,
Republic of North Macedonia 15–22

358. Ivan Boev, Dalibor Serafimovski, Goran Tasev

Geochemistry of recent sediments at confluence of the Blaštica river
into the Tikveš Lake 23–38

359. Trajče Stafilov, Robert Šajn, Mila Arapčeska, Ivan Kungulovski, Jasminka Alijagić

Multi-element distribution in road and street dust in the Bitola region,
North Macedonia 39–50

360. Aleksandra Trenchovska

Geomatics and GIS based system for geodiversity inventory and evaluation
applied in Kratovo 51–66

361. Lazar Gjorgjiev, Todor Serafimovski, Kristian Jovanov

3D Modeling of the Borov Dol porphyry copper deposit,
Republic of North Macedonia 67–82

Instructions to authors 83–84

<i>Geologica Macedonica</i>	Vol.	No	pp.	Štip
<i>Geologica Macedonica</i>	Год.	Број	стр.	Штип
	34	1	1–84	2020

СОДРЖИНА

356. Миха Буклески

Инфрацрвена огледална рефлексiona спектроскопија на поликристални пресувани таблети од алуми – Споредба со спектрите на монокристали 5–14

357. Тена Шијакова-Иванова, Блажо Боев, Лидија Робева Чуковска

Прелиминарни истражувања на гетит и монацит од Сивец, Република Северна Македонија 15–22

358. Иван Боев, Далибор Серафимовски, Горан Тасев

Геохемија на современите седименти на вливот на реката Блаштица во Тиквешко Езеро 23–38

359. Трајче Стафилов, Robert Šajn, Мила Арапческа, Иван Кунгуловски, Jasminka Alijagić

Мултиелементна дистрибуција на тешки метали во правот од улиците и патиштата во регионот на Битола, Северна Македонија..... 39–50

360. Александра Тренчовска

Инвентаризација и евалуација на геодиверзитетот во Кратово со примена на географски информациски системи и геоматика 51–66

361. Лазар Ѓорѓиев, Тодор Серафимовски, Кристијан Јованов

Тродимензијално моделирање на бакарното порфирско наоѓалиште Боров Дол, Република Северна Македонија..... 67–82

Упатство за авторите..... 83–84

PRELIMINARY INVESTIGATIONS OF GOETHITE AND MONAZITE FROM SIVEC, REPUBLIC OF NORTH MACEDONIA

Tena Šijakova-Ivanova¹, Blažo Boev¹, Lidija Robeva Čukovska²

¹*Faculty of Natural and Technical Sciences, Institute of Geology, “Goce Delčev” University in Štip,
Blvd. “Krstе Misirkov”, P.O. box 201, 2001 Štip, Republic of North Macedonia*

²*National Conservation Centre – Skopje. St. Josif Mihajlović 7, 1000 Skopje, Republic of North Macedonia
tena.ivanova@ugd.edu.mk*

A b s t r a c t: This paper presents the preliminary investigations of goethite and monazite that appear in marbles from Sivec. The presence of these minerals was not previously reported in the Sivec marbles that, along with the dominant dolomite and calcite, also revealed the presence of a vast number of other minerals including fluorite, rutile, phlogopite, corundum, diasporite, almandine, clinocllore, muscovite, quartz and zoisite. Goethite [Fe³⁺O(OH)] from Sivec occurs in small crystals found generally in fine-grained parts of the white dolomitic marble. The color ranges from yellow-brown to nearly black, and the luster varies correspondingly from earthy to submetallic. Cleavage is perfect {010} and less perfect {100}. The streak is yellowish brown. The size of the goethite crystals is 1 cm. The crystal system is orthorhombic. *Point Group:* 2/m 2/m 2/m, *Space Group:* Pbnm. The *unit cell* parameters obtained using the strongest X-ray diffraction peaks are: $a = 4.617 \text{ \AA}$, $b = 9.976 \text{ \AA}$, $c = 3.024 \text{ \AA}$, $Z = 4$, $V = 139.32 \text{ \AA}^3$. Monazite crystals are found on the surface of the goethite crystal. The size of the crystals within size varies in length (21.88 – 53.61 μm) and width (8.45 – 20.38 μm).

Key words: Sivec; marble; goethite; monazite

INTRODUCTION

The marbles mass from Sivec is a part of the Precambrian Pelagonia marble series, which is preserved along the eastern peripheral part of the Pelagonia crystalline mass. The geological setting of the immediate vicinity of Sivec includes the following members: Neogene and Quaternary sediments; the Riphean Cambrian metamorphic complex (phyl-lithoids, metadiabases, amphibole schists and marbles); granitoids; marble series: a) dolomites and calcithecical marbles, b) calcite marbles; mixed series (albite gneisses, micaschists, marbles and cipolins); series of gneissic micaschists (gneisses, mica-schists, quartzites and amphibolites) (Figure 1). The main lithostratigraphic features of Pelagonia derive from the primary accumulation of pelitic-psamitic and carbonate sediments accompanied by poorly expressed initial magmatism. In general, the lowest parts of the marble mass in Sivec consist of white dolomitic marbles of a granolastic composition and massive texture. According to the mineralogy,

structural and textural composition, the marble mass in Sivec is made of sucrose white dolomitic marble, dolomitic marble with planes, nests and calcite, and finely crafted calcite marble. Their dolomite composition is partly related to the primary gneiss. The structural features of the marbles from Sivec are reported by Dumurdžanov and Stefkov (1994).

The Sivec marble mass has exceptional, unique petrographic and mineralogical characteristics in terms of their structure and present mineral association that is a consequence of the specific genesis of this part of the Pelagonia marble metamorphic complex. Dolomite grain size ranges from 0.1 to 0.5 mm. The grains are partially rounded and exhibit pronounced crushing. Dolomite grains rarely have an irregular polygonal shape.

The influence of the granites that are near the marble mass is manifested by a higher presence of corundum, fluorite, paragonite, fangite, titanite, rutile, epidotic minerals, chlorite, diasporite and others.

The mineralogy of marbles from Sivec was previously investigated by several research groups: Erdmannsdorfer (1925), Barić (1960, 1969), Stojanov (1960, 1967, 1968), Jeršek and Mitrič (1999), (Jovanovski et al. (2012), Boev (2015), Šijakova-

Ivanova and Petreski (2018), Šijakova-Ivanova and Robeva Čukovska (2019). In all of the mentioned papers, the presence of goethite and monazite was not reported.




Fig. 1. Geological map of the area with the marble mass of Sivec-Pletvar-Kozjak (Dumurdžanov et al., 1990), R=1:18000. Legend: 1. Neogene and Quaternary sediments; 2. Riphean-Cambrian metamorphic complex (filithoids, metadiabases, amphibole shales and marbles); 3. granitoides; 4. marble series; 5. mixed series (albitic gneiss, micaschists, marbles and cypolines); 6. gneiss-micaschists series; 7. fault; 8. surface excavation of marbles

ANALYTICAL METHODS

For our research, the following analytical methods were used: Fourier transform (FT) infrared spectroscopy, scanning electron microscopy coupled with energy dispersive X-ray spectrometer, and X-ray powder diffraction.

Infrared spectroscopy. The FT infrared spectrum of the goethite sample was recorded on the IR-Prestige 21 spectrometer (Shimadzu, Japan) using the KBr pellet method, in the spectral range of 400–4000 cm^{-1} at a resolution of 2 cm^{-1} and 60 scans.

The pellet was prepared by mixing 0.3 mg of the sample powder and 100 mg of KBr. The spectra manipulation was performed by using the IR Solution 1.5 software (Shimadzu Corporation).

Scanning electron microscopy SEM analyses and electron micro-photographs were conducted using a VEGA3- LMU scanning electron microscopy. The study utilized semi-quantitative analysis using appropriate standards. The standards used are as follows: O: SiO_2 ; Na: albite; Mg: MgO; Al: Al_2O_3 ; Si:

SiO₂; P: GaP; Ca: wollastonite; Ti: Ti; Fe: Fe; Br: KBr. The results of the SEM/EDS analyses of mineral phases demonstrated the adequacy of this method for identification and characterization of mineral phases whose size is often below the resolution of an optical microscope.

XRD analyses were carried out by conventional X-ray diffraction techniques using the (Shimadzu) XRD-6100 diffractometer with CuK α (1.54178)

radiation operating at 40 kV and 30 mA. The powdered sample was scanned over the 10–80° range with a step size of 0.02° and scanning speed of 1.2°/min. The most intense registered maxima in the studied powder diagrams were compared with the corresponding diagrams from the ICDD PDF-2 software (2019). The Unit Cell software (Holland & Redfern, update 16th April 2006) was used for calculation on unit cell data.

RESULTS AND DISCUSSION

The goethite [Fe³⁺O(OH)] from Sivec occurs in small crystals found generally in fine-grained parts of the white dolomitic marble. The color ranges from light brown to nearly black and the luster varies from earthy to submetallic (Figure 2). The cleavage is perfect {010} and less perfect {100}. The streak is yellowish brown. The size of the goethite crystals is 1 cm.



Fig. 2. Crystal of goethite (1 cm length and 1cm width)

Showing the IR spectra characterization of limonite minerals, Makreski et al. (2004) revealed the occurrence of goethite (α -FeOOH) in the Allchar site and lepidocrocite (γ -FeOOH) from Pehčevo.

The IR spectrum of the examined sample of goethite is shown in Figure 3. The obtained data together with the corresponding literature data are shown in Table 1.




Fig. 3. Infrared spectrum of the goethite sample

Table 1

IR data of the examined sample from Sivec, compared with literature data

This work	Makreski et. al., 2004	Chukanov, 2014	Ruan et. al., 2002	
Goethite (Sivec)	Goethite (Allchar)	Goethite (Attica, Greece)	Goethite (Unheated)	
3417	3419	3300sh	3479	ν (H-O-H)
3155	3117	3120s	3206	ν (OH)
1646	1637	1655w	1686	δ (O-H)
1617			1637	γ (O-H)
1094	1096	–	–	–
1022	1017	1030sh	–	–
890	893	894	890	δ (OH)
798	799	797	799	γ (OH)
–	667	655sh	–	–
572	570	595	–	–
460*	472	465s	–	–
417*	407	412s	–	–

* Ascribed to ν (Fe-O...Fe) by Plyusnina (1977), cited by Makreski et. al., 2004

The most obvious spectral feature is the appearance of two well-separated bands at 890 and 798 cm^{-1} attributed to the hydroxyl deformation modes in the goethite structure. In the hydroxyl stretching region, two prominent bands at 3417 and 3155 cm^{-1} are observed. The band at 3155 cm^{-1} is located in a sub-region attributable to the stretching vibration of O-H units in goethite (Ruan et al., 2002). The band at 3417 cm^{-1} in line with the bands at 1646 and 1617 cm^{-1} can be assigned to the $\nu(\text{H}-\text{O}-\text{H})$ and $\delta(\text{H}-\text{O}-\text{H})$, respectively. These bands are indicative for the presence of non-stoichiometric hydroxyl units in the goethite structure (Ruan et al., 2002). Furthermore, the IR spectrum shows two less intensive distinct bands at 1096 and 1022 cm^{-1} which could be related to the characteristic modes of other minerals. Thus, it can be assumed that a small amount of additional mineral phase is also present in the goethite sample. The lower wave-number region exhibits an intense band with two separated maxima (460 and 417 cm^{-1}) and shoulder at 570 cm^{-1} which, according to Plyusnina (1977) and (Makreski et. al. (2004), are associated with $\nu(\text{Fe}-\text{O})$.

The results of the chemical analyses for the examined goethite and the comparison with literature data (Deer et al., 1962) are given in Figure 4 and Table 2.

The presence of Cu in the examined sample was determined, which can be seen from Table 2. The incorporation of Cu in the structure of natural goethite is more surprising because this element has a lower geochemical affinity for Fe oxides and a higher affinity for Mn oxides and phyllosilicates (Manceau et al., 2000). Other elements, which are commonly associated in goethite, such as Al, Si, P and Mn, were not detected in the goethite sample from Sivec.

On the surface of the goethite crystal, small monazite crystals were identified. Monazite is a rare earth phosphate mineral $\text{REE}(\text{PO}_4)$ containing mostly light REE and some heavy REE, more than bastnäsité (Kanazawa and Kamitani, 2006).

The generic chemical formula for monazite, $(\text{Ce}, \text{La}, \text{Nd}, \text{Th})(\text{PO}_4, \text{SiO}_4)$, reveals that cerium, lanthanum, neodymium and thorium can substitute for one another in the mineral's structure, and substitution of silica for phosphate also occurs.

According to George Blankson (Abaka-Wood et al., 2019), the REE minerals (bastnäsité, monazite, florencite, stillwellite, brannerite and stetindite) identified were in intimate associations with weakly ferromagnetic/strongly paramagnetic iron oxides (mainly hematite and goethite) and diamagnetic silicate (quartz, illite and annite) minerals.

The results of chemical analyses for monazite and comparison with literature data (Palache et al., 1951) are given in Figure 5 and Table 3.

The size of the monazite crystals varies in length 21.88 – 53.61 μm and width 8.45 – 20.38 μm .

Figure 6 represents the XRPD pattern of the investigated goethite.


The most intense registered maxima in the studied X-ray powder diagram were compared with the corresponding maxima in the diagram of goethite, ICDD 00 017 0536. The strongest peak is obtained at 2θ around 21° . This result is in compliance with that of Villacís-García et al. (2015), which reported goethite exhibits the highest peak intensity at $2\theta(21^\circ)$.

The unit cell parameters obtained using the main reflection lines of X-ray diffraction are $a = 4.617 \text{ \AA}$, $b = 9.976 \text{ \AA}$, $c = 3.024 \text{ \AA}$, $Z = 4$, $V = 139.32 \text{ \AA}^3$. The crystal system is orthorhombic. *Point Group*: 2/m 2/m 2/m, *Space Group*: Pbnm.


Table 2

Chemical composition of goethite from Sivec by SEM-EDX (weight %)


Element	Goethite from Sivec				Goethite from El Paso Co., Colorado, USA. Deer, et al., (1962)
	Analysis				
	1	2	3	4	
O	39.00	39.45	39.80	36.56	36.01
Fe	61.00	59.53	60.20	63.44	62.85
Cu	—	1.01	—	—	—
H	—	—	—	—	1.13
Total	100.00	100.00	100.00	100.00	100.00




a) SEM image of goethite (analysis 1)




b) EDX spectrum of goethite (analysis 1)




c) SEM image of goethite (analysis 2)




d) EDX spectrum of goethite (analysis 2)




e) SEM image of goethite (analysis 3)



f) EDX spectrum of goethite (analysis 3)




g) SEM image of goethite (analysis 4)




h) EDX spectrum of goethite (analysis 4)


Fig. 4. SEM image and EDX spectrum of goethite from Sivec




a) SEM image of monazite (analysis 1)




b) EDX spectrum of monazite (analysis 1)




c) SEM image of monazite (analysis 2)




d) EDX spectrum of monazite (analysis 2)




e) SEM image of monazite (analysis 3)



f) EDX spectrum of monazite (analysis 3)



g) SEM image of monazite (analysis 4)



h) EDX spectrum of monazite (analysis 4)

Fig. 5. SEM image of monazite and EDX spectrum of monazite from Sivec

Table 3

Chemical composition of monazite from Sivec by SEM-EDX (weight %)

Element	Monazite from Sivec				Monazite from Mars Hill, Madison County, North Carolina (Palache et al., 1951)
	1	2	3	4	
O	31.30	32.92	28.66	40.05	26.64
P	14.01	14.05	12.56	13.58	12.89
La	16.16	15.80	18.39	14.80	14.46
Ce	27.87	27.13	31.76	24.38	29.17
Nd	9.67	10.10	8.63	7.20	12.01
F	0.99	—	—	—	—
Total	100.00	100.00	100.00	100.00	100.00




Fig. 6. XRD pattern of the investigated goethite from Sivec

CONCLUSION

This study describes the occurrence of goethite and monazite in the marble mass from Sivec for the first time. Goethite occurs in single crystals that occurred from the alteration of pyrite and it is found generally in the fine-grained parts of the white dolomitic marble. Pseudomorph of goethite has completely replaced pyrite from the matrix. The oxidation of iron sulfides invariably leads to the formation of

goethite [α -FeO(OH)] in intimate association with the sulfides. The size of the goethite crystals is 1 cm. The color ranges from yellow – brown to nearly black. On the surface of the goethite crystal, small monazite crystals are found. The size of the monazite crystals varies in length 21.88 – 53.61 μm and width 8.45 – 20.38 μm .

REFERENCES

- Abaka-Wood, G. B., Zanin, M., Addai-Mensah J., Skinner, W. (2019): Recovery of rare earth elements minerals from iron oxide-silicate rich tailings – Part 1: Magnetic separation. *Minerals Engineering*, Vol. **136**, pp 50–61.
- Barić, L. (1960): Optische Eigenschaften des Diaspors von Sivec unweit von Prilep in Mazedonien, *Bulletin of the Scientific Council of Academy of Yugoslavia*, **5**, 71–76.
- Barić, L. (1969): Mineralogical discoveries that have so far been made at material sites in Macedonia, *Geological Survey of Macedonia, Book of Papers*, 14, 55–66 (in Macedonian).
- Boev, B. (2015): Sivec type of marble (mineralogical, whiteness and physico-mechanical features). *Geologica Macedonica*, **29** (2), pp. 167–176. ISSN 0352-1206.
- Chukanov, N. V., (2014): *Infrared Spectra of Mineral Species*, Extended Library, Vol. **1**, Springer, p. 331.
- Deer, W. A., Howie, R. A., Zussman, J. (1962): *Rock Forming Minerals*: Vol. **1–5**. Longmans, Green & Co., London.
- Dumurdžanov, N., Petrov, G. (1990): Lithostratigraphy and chemical characteristic on Vardar oceanic crust on territory of S.R. Macedonia. *XII Congress of Geologists of Yugoslavia*, Ohrid, pp. 51–63.
- Dumurdžanov, N., Stefkov, N. (1994): The geology and technical features of the Sivec marble deposit – Prilep, Republic of Macedonia, *Geologica Macedonica*, **8**, 29–35.
- Erdmannsdorfer, O. H. (1925): Über Kosmatite in neues Glied der Sprodglimmergruppe und seine Parageneses, *Zentralblatt für Mineralogie, Geologie, Paleontologie*, Abteilung A, 69–72.
- Griffiths, P. R., De Haseth, J. A., Winefordner, J. D. (Series Editor) (2007): *Fourier Transform Infrared Spectrometry*, Wiley, e-book, 2nd edition 560 p. ISBN: 978-0-471-19404-0.

- Holland, T., Redfern, S. (1997,) new version April 2006): Unit cell refinement from powder diffraction data: the use of regression diagnostics. *Mineralogical Magazine*, 61, 65–77.
- ICDD PDF-2 (2019): International Centre for Diffraction Data, Softwer program.
- Jeršek, M., Mitrič, B. (1999): Corundum from Prilep Dolomitic Marble (Macedonia), *Scopolia, Journal of the Slovenian Museum of Natural History, Ljubljana*, No. 41, September 1999, pp. 1–22.
- Jovanovski, G., Boev, B., Makreski, P. (2012): *Minerals from the Republic of Macedonia with Introduction to Mineralogy*, MANU, Skopje.
- Kanazawa, J., Kamitani, M. (2006): Rare earth minerals and resources in the World, *Journal of Alloys and Compounds*, 408–412 (19), pp. 1339–1343.
- Makreski, P., Jovanovski, G., Kaitner, B., Stafilov, T., Boev, B., Cibrev, D. (2004): Minerals from Macedonia. X. Separation and identification of some oxide minerals by FTIR spectroscopy, AAS, AES-ICP and Powder XRD, *Neues Jahrb. Mineral. Abh.*, **180**, pp. 215–243.
- Manceau, A., Schlegel, M. L., Musso, M., Sole, V. A., Gauthier, C., Petit, P. E. (2000): Crystal chemistry of trace elements in natural and synthetic goethite, *Geochimica et Cosmochimica Acta*, Vol. **64**, No. 21, pp. 3643–3661.
- Palache, C., Berman, H., Frondel, C. (1951): *The System of Mineralogy of James Dwight Dana and Edward Salisbury Dana*, Yale University 1837–1892, Volume II: 695.
- Plyusnina, I. I. (1977): *Infrakrasnye spektry mineralov*, Moskva. Izd. Moskovskogo Universiteta (*Infrared Spectra of Minerals*, Moscow).
- Ruan, H. D., Frost, R. L., Klopogge, J. T., Duong, L. (2002): Infrared spectroscopy of goethite dehydroxylation: III. FT-IR microscopy of in situ study of the thermal transformation of goethite to hematite, *Spectrochimica Acta, Part A*, 58, pp. 967–981.
- Stojanov, R. (1960): Previous results from geological and petrological research on high metamorphic rocks in the central part of the Pelagonia massif. Excavations of high metamorphic rocks in the central part of the Pelagonia massif. *Geological Institute Papers* 13, Skopje.
- Stojanov, R. (1967): *Petrological characteristics of magmatic and metamorphic rocks from the wider area of Prilep*, Doctoral dissertation, Faculty of Mining and Geology, Beograd.
- Stojanov R. (1968): Fengites in Pelagonian metamorphic complex, Geological Survey of Macedonia, *Book of Papers*, **13**, 41–57 (in Macedonian).
- Šijakova-Ivanova, T., Petreski, M. (2018): Identification of minerals from Sivec with application of XRD method. *Podex-Poveks '18*, **XI**, pp. 244–253. ISSN ISBN 978-608-65530-4-3
- Šijakova-Ivanova, T.; Robeva Čukovska, L. (2019): Preliminary investigations of paragonite in marbles from Sivec, North Macedonia *Geologica Macedonica*, Vol. **33**, No. 1, pp. 53–59.
- Villacís-García, M., Ugalde-Arzate, M., Vaca-Escobar, K., Villalobos, M., Zanella, R., Martínez-Villegas, N. (2015): Laboratory synthesis of goethite and ferrihydrite of controlled particle sizes, *Boletín de la Sociedad Geológica Mexicana*, Vol. **67**, No. 3, pp. 433–446. .

Резиме

ПРЕЛИМИНАРНИ ИСТРАЖУВАЊА НА ГЕТИТ И МОНАЦИТ ОД СИВЕЦ, РЕПУБЛИКА СЕВЕРНА МАКЕДОНИЈА

Тена Шијакова-Иванова¹, Блажо Боев¹, Лидија Робева Чуковска²

¹Институтот за Геологија, Факултетот за природни и технички науки, Универзитетот „Гоце Делчев“ во Штип,
бул. „Крсте Мисирков“, б.ф.п. 201, 2001 Штип, Република Северна Македонија

²Национален центар за конзервација, Скопје, Република Северна Македонија
tena.ivanova@ugd.edu.mk

Клучни зборови: Сивец; мермер; гетит; монацит

Во овој труд се презентираат резултати од прелиминарните истражувања на кристали на гетит кој се појавува во мермерите од Сивец. Испитаниот примерок беше анализиран со методите IR, SEM-EDS и XRD. Гетитот се јавува во мали кристали псевдоморфи по пирит, кои се наоѓаат главно во фино зрнести делови на белиот доломитски мермер. Кристалите се јавуваат поединечно. Бојата се движи од светло-кафеава до речиси црна, а сјајноста варира од земјена до полуметалична. Цепливоста е совршена по {010} и слабо совршена по {100}. Бојата на огребот е жолтеникаво кафеава. Големината на кристалот е 1 cm. Добиените податоци од рендгенските дифракциони испитувања

се споредени со соодветните максимуми од дијаграмот на гетит, ICDD 00 017 0536. Со добиените податоци од рендгенските дифракциони испитувања се пресметани димензиите на елементарната ќелија. Добиени се следниве резултати: $a = 4,617 \text{ \AA}$, $b = 9,976 \text{ \AA}$, $c = 3,024 \text{ \AA}$, $Z = 4$, $V = 139,32 \text{ \AA}^3$. Со скенирачка електронска микроскопија беше утврдено и присуство на кристали на монацит. Кристалите на монацит се наоѓаат на површината на кристалот на гетитот. Големината на кристалите на монацит варира во должина (21,88 – 53,61 μm) и ширина (8,45 – 20,38 μm). Оваа студија за првпат ја опишува појавата на гетит и монацит во мермерната маса од Сивец.